MANAGEMENT OF THE SICK YOUNG INFANT AGE UPTO 2 MONTHS FOR HW

Name:	Ag	ge: Gender:	Weight:kg Ten	nperature:°C / °F	Date:
ASK: What are	the infant's problen	18?	Initial visit?	_ Follow up visit?	
ASSESS (Circle	•				CLASSIFY
Is the infant ha	OSSIBLE SERIOUS B aving difficulty in feeding thad convulsions?		ION/JAUNDICE		
That the finance	nad conversions.		Count the oreating in one minute	eathing? ible, feel for fever or low °C (95.9°F/99.5°F) or e him/her mulated but then stops?	
Ask when did jaundice appeared – First 24 hours After 24 hours			 Look for skin pustules Look for jaundice (yellow skin), If present Look at the young infant's palms and soles. Are they yellow? 		
DOES THE YO	UNG INFANT HAS DI	ARRHOEA?	Yes No	<u></u>	
		•	Look at the young infant's general conditi Look at infant's movements: Does the infant move only when stin Does the infant not move at all? Is the infant restless and irritable? Look for sunken eyes. Pinch the skin of the abdomen. Does it go Slowly? Very slowly (longer than 2 seconds)	mulated and then stops? b back:	
THEN CHECK	FOR FEEDING PROI	BLEM & VERY LOW	WEIGHT		
• Does the indirinks? ⇒ If yes, ⇒ What d	ant breastfed in the previous times in 24 hours? times ant usually receive any Yes No how often? lo you use to feed the interactions for urgent References.	other foods or	YesNo		
 If infant ha ask the mot 	STFEEDING: ant breastfed in the previous not breastfed in the previbent to put her infant to the breastfed for 4 minutes.	ious hour? evious hour, he breast.	 ⇒ Mouth wide open Yes_ ⇒ Lower lip turned outward Yes_ ⇒ More areola above than below 	No, slow deep sucks, feeding?	
CHECK THE Y					Circle immunization needed today Return for next immunization on:
Birth 6 weeks	BCG Penta-1	OPV-1	HEP-B 0 Rotavirus-1 flPV-1	PCV-1	(Date)
ASSESS CAREO ASK: How do you play How do you talk! How do you get y	with your baby? to your baby? /our baby smile?	S TO SUPPORT CHII	Look how does caregiver show he/she is movement? Look how does caregiver comfort the ch		

•	Advise Mother to Give Home Care to the Young Infant	
	Immediately after birth, baby should be put on the mother's	
	abdomen for skin to skin contact.	
	Initiate breastfeeding within one hour of birth.	
	➤ Breastfeed day and night as often as your baby wants, at least 8	
	times in 24 hours. Frequent feeding produces more milk.	
	➤ If your baby is small (low birth weight), feed him or her at least every 2-3 hours. Wake the baby for feeding after 3 hours, if she	
	or he does not wake self.	
	Breastfed as often as your child wants. Look for signs of hunger,	
	such as beginning to fuss, sucking fingers, or moving lips.	
	DO NOT give other foods or fluids. Breast milk is all your baby	
	needs.	
	Make sure the young infant stays warm at all times. In cool	
	weather, cover the infant's head and feet and dress the infant	
	with extra clothing.	
	Advise mother to wash hands with soap and water after defecation and after cleaning bottom of the baby	
	Do not apply anything on the cord and keep the umbilical cord	
	dry.	
	-	
•	Advise the mother to return immediately if the young infant has	
	any of these danger signs:	
	 Not able to drink or breastfeed Becomes sicker 	
	Develops a fever or feels cold to touch	
	Fast breathing	
	> Difficult breathing	
	Yellow palms and soles (if infant has jaundice)	
	> Blood in stool	
	-	
	-	
	-	Counsel mother shout feeding
		Counsel mother about feeding
		Counsel mother about development supportive practices
		2 mome about acrosophical supportate practices
		Advise mother when to return immediately.
		*
		Cive any immunization needed today
		Give any immunization needed today
		Counsel the mother about her own health
		Return to follow up in: